

myProbes Passport (RUO Probes)

MPD4470

IGH/BCL6 Dual Fusion Probe

For Research Use Only

Nucleotide Locations (If requesting more than one probe please specify individual probe colours)

	Chromosome Locus	Start	End	Colour
1	3q27.3	187061090	187209930	Red
2	3q27.3	187211778	187376421	Red
3	3q27.3	187331758	187475894	Red
4	3q27.3	187446099	187594575	Red
5	3q27.3	187624268	187753558	Red
6	14q32.33	105639058	105802170	Green
7	14q32.33	105788920	105998428	Green
8	14q32.33	106667215	106795977	Green
9	14q32.33	106782750	106874172	Green
10	14q32.33	106841162	106991516	Green
11	14q32.33	106995083	107135808	Green
12	14q32.33	107077128	107174325	Green
13	14q32.33	107160170	107267431	Green
14	14q32.33	107258048	107284021	Green


Database Used and Version


GRCh37 (hg19)





Probe Map





Materials Provided

1. **Probe:** 100μl per vial

The probe is provided pre-mixed in hybridisation solution (Formamide; Dextran Sulphate; SSC) and is ready to use. It is directly labelled with a red fluorophore, which lies in the Texas Red Spectrum and a green fluorophore, which lies in the FITC Spectrum. The probe has been batch released after QC testing on Formalin Fixed Paraffin Embedded Sample and Bone Marrow Sample.

2. **Counterstain:** 150μl per vial. The counterstain is DAPI antifade (ES: 0.125μg/ml DAPI (4,6-diamidino-2-phenylindole)).





Warnings and Precautions

- 1. For research use only. Not for use in diagnostic procedures.
- 2. For laboratory professional use only.
- 3. Probe mixtures contain formamide, which is a teratogen; do not breathe fumes or allow skin contact. Handle with care; wear gloves and a lab coat.
- 4. DAPI is a potential carcinogen. Handle with care; wear gloves and a lab coat.
- 5. Follow local disposal regulations for your location along with recommendations in the Safety Data Sheet to determine the safe disposal of this product. This also applies to damaged test kit contents.
- 6. Dispose of all used reagents and any other contaminated disposable materials following procedures for infectious or potentially infectious waste. It is the responsibility of each laboratory to handle solid and liquid waste according to their nature and degree of hazardousness and to treat and dispose of them (or have them treated and disposed of) in accordance with any applicable regulations.
- 7. Operators must be capable of distinguishing the colours red, blue, and green.
- 8. The probe should not be diluted or mixed with other probes.
- 9. All products should be validated before use.
- 10. Internal controls should be carried out by using unaffected cell populations in testing samples.
- 11. Custom probes are specifically developed for individual customers' RESEARCH USE ONLY (RUO) requirements and not with the intention of being used for in vitro diagnostic examination. Therefore, prior to any use of these probes, users should review the design of such probes to confirm they are suitable for their requirements.

Storage and Handling

- 1. Store the probe between -25°C to -15°C.
- 2. Based on the stability established for other substantially equivalent CytoCell probes, this myProbes product should be stable for 2 years post manufacture date when stored as indicated on the label.
- 3. Store the probe and counterstain vials in the dark. Ensure that exposure of the probe and counterstain to laboratory lights is limited at all times.

Known Cross-Reactivity

The IGH probe may show faint cross hybridisation to a region on Chromosome 15 and 16.

Customer Support

Please contact the CytoCell Technical Support Department or email probes@cytocell.com.

Patents and Trademarks

myProbes and CytoCell are registered trademarks of Cytocell Ltd.

For Research Use Only. Not for use in diagnostic procedures.

This product contains technology licensed from Life Technologies Corporation that is available for human diagnostics or life science research use only.



Cytocell Ltd.

Oxford Gene Technology 418 Cambridge Science Park, Milton Road,

Cambridge, CB4 0PZ, UK T: +44(0)1223 294048

F: +44(0)1223 294986

E: probes@cytocell.com

W: www.ogt.com

